

# 2011 Air Quality Monitoring Data Summary

---



**State of Idaho  
Department of Environmental Quality**

**April 2015**



Printed on recycled paper, DEQ, April 2015, MPOB, 81630. Costs associated with this publication are available from the State of Idaho Department of Environmental Quality in accordance with Section 60-202, Idaho Code.

# **2011 Air Quality Monitoring Data Summary**

**April 2015**



**Idaho Department of Environmental Quality  
Air Quality Division  
1410 N. Hilton  
Boise, ID 83706**

The 2011 Air Quality Monitoring Data Summary is available for viewing or downloading on the DEQ website at <http://www.deq.idaho.gov/air-monitoring-network>.

Links to additional documents for download are also available at the DEQ website.

## Table of Contents

1	Introduction.....	1
2	Air Quality Standards .....	3
3	Monitoring Network .....	5
3.1	Pollutant Monitoring .....	8
4	Monitoring Results .....	9
4.1	Ozone.....	9
4.2	Particulate Matter (10 micrometers).....	20
4.3	Particulate Matter (2.5 micrometers).....	22
4.4	Carbon Monoxide.....	44
4.5	Sulfur Dioxide .....	45
4.6	Lead .....	46
4.7	Nitrogen Dioxide .....	47
5	Air Quality Index.....	49
6	Impaired Air Quality.....	57
6.1	Winter Burn Bans .....	57
6.2	Summer Ozone Alerts .....	57
7	Definitions .....	58
8	Pollution Sources .....	59
9	Criteria Air Pollutants.....	60
	Air Quality Data Summaries.....	65

## List of Figures

Figure 1.	2011 Idaho ambient air monitoring network.....	7
Figure 2.	2011 Idaho nonattainment and maintenance areas.....	11
Figure 3.	Coeur d'Alene—Lancaster Road 8-hour ozone maximum concentration.....	12
Figure 4.	Coeur d'Alene—Lancaster Road highest 8-hour ozone concentrations and 3-year average of the 4th highest concentration.....	13
Figure 5.	Meridian—St. Luke's 8-hour ozone daily maximum concentration.....	14
Figure 6.	Meridian—St. Luke's highest 8-hour ozone concentrations and 3-year average of the 4th highest concentration.....	15
Figure 7.	Boise—Idaho Transportation Department 8-hour ozone daily maximum concentration.....	16
Figure 8.	Boise—Idaho Transportation Department annual four highest 8-hour ozone concentrations and 3-year average of the 4th highest concentration.....	17
Figure 9.	Boise—White Pine 8-hour ozone daily maximum concentration.....	18
Figure 10.	Boise—White Pine highest 8-hour ozone concentrations and 3-year average of the 4th highest concentration.....	19
Figure 11.	Idaho 8-hour ozone concentrations and 3-year average of the 4th highest concentration.....	20

Figure 12. Three-year average of daily maximum PM<sub>10</sub>. ..... 21

Figure 13. Three-year average estimated exceedances of the daily PM<sub>10</sub> standard..... 22

Figure 14. Three-year average 98th percentile daily PM<sub>2.5</sub> concentration (monitors operated in 2011)..... 24

Figure 15. PM<sub>2.5</sub> 3-year average annual mean (monitors operated in 2011). ..... 25

Figure 16. Nampa—Fire Station PM<sub>2.5</sub> daily averages from continuous analyzer..... 26

Figure 17. Coeur d'Alene—Lancaster Road PM<sub>2.5</sub> daily averages from continuous analyzer. .... 27

Figure 18. St. Maries PM<sub>2.5</sub> daily averages from continuous analyzer..... 28

Figure 19. Sandpoint PM<sub>2.5</sub> daily averages from continuous analyzer. .... 29

Figure 20. Lewiston PM<sub>2.5</sub> daily averages from continuous analyzer. .... 30

Figure 21. Moscow PM<sub>2.5</sub> daily averages from continuous analyzer. .... 31

Figure 22. Twin Falls PM<sub>2.5</sub> daily averages from continuous analyzer. .... 32

Figure 23. Pocatello—Garrett and Gould PM<sub>2.5</sub> daily averages from continuous analyzer. .... 33

Figure 24. Idaho Falls PM<sub>2.5</sub> daily averages from continuous analyzer. .... 34

Figure 25. Garden Valley PM<sub>2.5</sub> daily averages from continuous analyzer..... 35

Figure 26. Grangeville PM<sub>2.5</sub> daily averages from continuous analyzer. .... 36

Figure 27. Idaho City PM<sub>2.5</sub> daily averages from continuous analyzer. .... 37

Figure 28. Ketchum PM<sub>2.5</sub> daily averages from continuous analyzer. .... 38

Figure 29. McCall PM<sub>2.5</sub> daily averages from continuous analyzer. .... 39

Figure 30. Meridian—St. Luke's PM<sub>2.5</sub> daily averages from continuous analyzer..... 40

Figure 31. Franklin PM<sub>2.5</sub> daily averages from continuous analyzer..... 41

Figure 32. Pinehurst PM<sub>2.5</sub> daily averages from continuous analyzer. .... 42

Figure 33. Salmon PM<sub>2.5</sub> daily averages from continuous analyzer..... 43

Figure 34. Carbon monoxide 2nd highest 8-hour concentration. .... 45

Figure 35. Nitrogen dioxide annual 1-hour average. .... 48

Figure 36. Air quality for Ada County..... 51

Figure 37. Air quality for Bannock County..... 51

Figure 38. Air quality for Benewah County. .... 51

Figure 39. Air quality for Bonner County. .... 52

Figure 40. Air quality for Butte County..... 52

Figure 41. Air quality for Canyon County..... 52

Figure 42. Air quality for Caribou County. .... 53

Figure 43. Air quality for Cassia County..... 53

Figure 44. Air quality for Custer County..... 53

Figure 45. Air quality for Franklin County..... 54

Figure 46. Air quality for Idaho County. .... 54

Figure 47. Air quality for Kootenai County..... 54

Figure 48. Air quality for Latah County. .... 55

Figure 49. Air quality for Lemhi County..... 55

Figure 50. Air quality for Nez Perce County..... 55

Figure 51. Air quality for Shoshone County..... 56

Figure 52. Air quality for Twin Falls County. .... 56

## List of Tables

Table 1. 2011 air quality standards for criteria pollutants. ....	4
Table 2. Monitoring network for 2011. ....	6
Table 3. Monitoring methods used in Idaho in 2011. ....	8
Table 4. US Environmental Protection Agency Air Quality Index breakpoint definitions. ....	49
Table 5. 2011 Air Quality Index yearly summary. ....	50

*This page intentionally left blank for correct double-sided printing.*

## 1 Introduction

This annual report is issued by the Idaho Department of Environmental Quality (DEQ) to inform the public of air quality throughout Idaho. The purpose of this report is to summarize regional ambient air quality while presenting air monitoring results for six criteria air pollutants. The US Environmental Protection Agency (EPA) sets national ambient air quality standards (NAAQS) for these pollutants. These criteria air pollutants include the following:

- Particulate matter ( $PM_{10} \leq 10$  micrometers [ $\mu m$ ] and  $PM_{2.5} \leq 2.5 \mu m$  in diameter)
- Carbon monoxide (CO)
- Sulfur dioxide (SO<sub>2</sub>)
- Nitrogen dioxide (NO<sub>2</sub>)
- Ozone (O<sub>3</sub>)
- Lead (Pb)

In Idaho, criteria pollutant monitoring occurs primarily in areas of high population where the potential for human exposure is greatest. Particulate matter is currently the most common criteria air pollutant of concern in Idaho because particulate sources are widespread throughout the state. Common sources include windblown dust, reentrained road dust, smoke (residential, agricultural, and forest fires), industrial emissions, and motor vehicle emissions.

The  $PM_{10}$  standard has been in effect since 1987 and historically had been the particulate size of concern. However,  $PM_{2.5}$ , or fine particulate matter, has been monitored in Idaho since 1998 and has become a pollutant of concern. Numerous studies have associated  $PM_{2.5}$  with a variety of respiratory and cardiovascular problems, ranging from aggravated asthma to irregular heartbeats, heart attacks, and early death in people with heart or lung disease. The  $PM_{2.5}$  and  $PM_{10}$  NAAQS were revised by EPA effective December 17, 2006. Due to a lack of evidence linking health problems to long-term exposure to coarse particle pollution, EPA revoked the annual  $PM_{10}$  standard of 50 micrograms per cubic meter ( $\mu g/m^3$ ) while retaining the short-term 24-hour standard of 150  $\mu g/m^3$ . The 24-hour standard for  $PM_{2.5}$  was lowered from 65  $\mu g/m^3$  to 35  $\mu g/m^3$  to provide increased protection against health effects associated with short-term exposure (including premature mortality and increased hospital admissions and emergency room visits).

Another historical air pollutant of concern in Idaho is carbon monoxide. The primary source of carbon monoxide is incomplete fossil fuel combustion. Carbon monoxide concentrations have the potential to be high in the urbanized areas where automobile traffic is heavy and cars frequently idle at stoplights. The Boise area (northern Ada County) was the only carbon monoxide nonattainment area in the state. When the State Implementation Plan (SIP) and Maintenance Plan were accepted by EPA on December 27, 2002, it was reclassified as a maintenance area. No violations of the 1- or 8-hour carbon monoxide NAAQS have occurred since 1991.

Sulfur dioxide and nitrogen dioxide sources are few and localized because these air pollutants come primarily from large industrial sources (transportation sources also contribute to nitrogen dioxide). Little heavy industry exists in Idaho, and elevated sulfur dioxide and nitrogen dioxide concentrations in ambient air are typically not found. However, due to potential concerns of some localized sources, DEQ has monitored for one or both of these pollutants in Boise,

Pocatello, Moyie Springs, Mountain Home, Coeur d'Alene, Meridian, and Soda Springs. In the past 10 years of targeted monitoring, DEQ has not measured significant concentrations of these pollutants at these monitoring sites.

The fifth criteria air pollutant, ozone, has been monitored by DEQ, in the Treasure Valley since 2002 and in Coeur d'Alene from 2005 to 2011. Ozone is created when combustion by-products (volatile organic compounds [VOCs]) near the ground react with nitrogen oxides and other compounds to create photochemical smog. These reactions are stimulated on days of intense sunlight and warm temperatures. Ozone has become a pollutant of concern since many summertime days are classified as moderate for ozone on the Air Quality Index (AQI). EPA lowered the 8-hour ozone standard on May 27, 2008, from 0.08 parts per million (ppm) to 0.075 ppm. The new standard poses a greater risk of nonattainment for all airsheds but particularly the Treasure Valley airshed. EPA announced it was reconsidering the ozone standard and was expected to release new proposed NAAQS ozone standards in December 2009. In 2011, EPA announced it would postpone any changes to the ozone NAAQS until 2013.

The sixth criteria air pollutant, lead, is not currently being monitored by DEQ. Lead was monitored in the Shoshone County town of Kellogg, near the Bunker Hill Superfund site, because lead was a by-product of the smelting process that occurred in the area for decades. Although a significant problem in the 1970s and early 1980s, airborne lead concentrations at this monitoring site were very low through the 1990s. DEQ discontinued monitoring for lead in 2002. EPA reviewed the lead NAAQS and on November 12, 2008, lowered the standard significantly to 0.15  $\mu\text{g}/\text{m}^3$ . The new standard provided different monitoring requirements based on whether sources were emitting significant volumes of lead. Source-oriented monitoring is required for states with sources of lead that emit or have the potential to emit more than 0.5 tons per year (tpy). Nonsource-oriented monitoring is required for urban areas with a population greater than 500,000. The nonsource-oriented requirements will be implemented at the NCore multipollutant monitoring station in Meridian. DEQ will initiate  $\text{PM}_{10}$  lead monitoring at the NCore site in Meridian by January 1, 2012.

The NCore multipollutant monitoring site in Meridian is part of an EPA network that uses advanced measurement systems to record data for particles, trace gases, and meteorology. The NCore data are used to support air quality forecasting and model evaluation, and to develop emissions strategies. The data are also used to assess compliance with the NAAQS.

DEQ monitored for certain common urban toxic air pollutants in the Treasure Valley from 2003 to the beginning of 2005 to determine if concentrations were at levels that could have adverse health effects. The Community Scale Air Toxics Monitoring Project also measured toxic air pollutants in 2007. Health effects from toxic air pollutants include, but are not limited to, increased cancer risk and respiratory, cardiovascular, and neurological effects. While DEQ has discontinued air toxics monitoring, the data proved valuable toward reconciling EPA's National Air Toxics Assessment (NATA) program to verify prediction models. The NATA models predict cancer and noncancer risk values across Idaho's airsheds using emissions estimates of certain air toxic compounds. The data have also been crucial in developing DEQ's air toxics models. As resources become available, Idaho may resume air toxic monitoring in the future.

While Idaho generally enjoys good air quality, in many ways our airsheds are faced with new challenges. Some of these challenges are related to long-term economic and population growth,

particularly in terms of vehicles on roadways and growth in new construction. Each day, DEQ measures the concentration of certain air pollutants throughout the state. DEQ may issue local burn restrictions (voluntary and/or mandatory) when concentrations of these air pollutants reach or exceed the health-based standards or limits established by local ordinance, state law, or federal regulation. Concerned citizens may tune in to the news on their local radio or television station to find out if a burn ban has been issued, or access DEQ's website at <http://www.deq.idaho.gov/air-quality.aspx>. DEQ issues a news bulletin to local news media, law enforcement, and fire officials each time a burn ban is imposed. Each year, a number of voluntary and sometimes mandatory bans are issued due to deteriorated local air quality conditions.

Real-time air monitoring data are available on DEQ's website at <http://airquality.deq.idaho.gov/>. We encourage you to visit our website at <http://www.deq.idaho.gov/> to find more extensive air quality data, educational materials, and discussions of current topics.

We are expanding and refining our website to better serve the residents of Idaho. Improvements are expected to provide the public with better access to real-time monitoring data as well as reorganize publications and other information regarding air quality. We want your feedback on our air quality data and program. Please submit your comments via e-mail to Bruce Louks, Modeling, Monitoring, and Emission Inventory Manager, at [Bruce.Louks@deq.idaho.gov](mailto:Bruce.Louks@deq.idaho.gov) or call at (208) 373-0294.

## **2 Air Quality Standards**

The federal Clean Air Act of 1970 (CAA) requires EPA to set NAAQS for air pollutants considered harmful to public health and the environment. The standards are designed to primarily protect the general public, including sensitive populations such as asthmatics, children, and the elderly. They are also intended to safeguard public welfare by reducing effects such as decreased visibility and damage to animals, crops, vegetation, and buildings. EPA established standards for six criteria air pollutants. Table 1 contains seven air pollutants, which include two size ranges of particulate matter.

Idaho adheres to the NAAQS. For more information, EPA air quality standards and supporting rationale are available at <http://epa.gov/air/criteria.html>.

**Table 1. 2011 air quality standards for criteria pollutants.**

Pollutant	Level	Averaging Time	Metric
<a href="#">Ozone (O<sub>3</sub>)</a>	0.075 ppm	8-hour	The 3-year average of the 4th-highest daily maximum 8-hour average concentration cannot exceed the level measured at each monitor within an area over each year. The standard was lowered May 27, 2008, from 0.08 ppm.
<a href="#">Particulate matter, 10 micrometers (PM<sub>10</sub>)</a>	150 µg/m <sup>3</sup>	24-hour	The 24-hour average cannot exceed the level more than once per year on average over 3 years.
<a href="#">Particulate matter, 2.5 micrometers (PM<sub>2.5</sub>)</a>	15 µg/m <sup>3</sup>	Annual (arithmetic average)	The 3-year annual average of the weighted annual mean concentrations cannot exceed the level. The standard was lowered December 17, 2006, from 15.4 µg/m <sup>3</sup> .
	35 µg/m <sup>3</sup>	24-hour	The 3-year average of the 98th percentile (based on the number of samples taken) of the daily concentrations must not exceed the level. The 24-hour standard was lowered from 65 µg/m <sup>3</sup> to 35 µg/m <sup>3</sup> on December 17, 2006.
<a href="#">Carbon monoxide (CO)</a>	35 ppm	1-hour	The 1-hour average cannot exceed the level more than once per year.
	9 ppm	8-hour	The 8-hour average cannot exceed the level more than once per year.
<a href="#">Sulfur dioxide (SO<sub>2</sub>)</a>	75 ppb	1-hour	The 99th percentile of 1-hour daily maximum concentrations, averaged over 3 years, cannot exceed the level. The 1-hour standard was revised in 2010.
<a href="#">Lead (Pb)</a>	0.15 µg/m <sup>3</sup>	Rolling 3-month average	The rolling 3-month average (12-average periods per year) cannot exceed the level. The standard was lowered October 15, 2008, from 1.5 µg/m <sup>3</sup> .
<a href="#">Nitrogen dioxide (NO<sub>2</sub>)</a>	0.053 ppm	Annual (arithmetic average)	The annual mean cannot exceed the level.
	100 ppb	1-hour	The 98th percentile of 1-hour values, averaged over 3 years cannot exceed the level.

Notes: Hyperlinks to pollutants are shown in blue. Daily concentration is the 24-hour average, measured from midnight to midnight. Parts per million (ppm); micrograms per cubic meter (µg/m<sup>3</sup>); parts per million (ppb)

The NAAQS for each pollutant may have different averaging periods (e.g., hourly and 8-hour averages). These different forms of the standard are created and enforced to address varied health impacts that result from shorter, high-level exposure versus longer, low-level exposure. These differences are addressed pollutant-by-pollutant in the following sections, and additional information is available on the EPA website. A distinction exists between *exceeding* and

*violating* a standard; the two are not equivalent. This distinction results from the nature of the standards. In most instances, it is allowable for an area to exceed the standard a few times to allow for possible unusual meteorological circumstances. For example, a carbon monoxide 8-hour average of 15 ppm clearly exceeds the standard; however, it does not violate the standard if it is the only exceedance that year (the standard allows for one exceedance).

The EPA standards typically apply to an *area*, which may be defined in different ways. Data are presented for individual monitoring stations in the following sections because this provides more insight into regional differences in Idaho's ambient air quality. The following summaries show how Idaho's airsheds compared to the standards discussed above for 2011 and in many instances incorporate the AQI and other measures of air quality where appropriate. The AQI color code shading is shown to aid in interpreting air quality but does not imply whether or not standards were met for each air pollutant. An airshed must satisfy the conditions in Table 1 to ensure compliance with the NAAQS.

### 3 Monitoring Network

The Idaho monitoring network is a composite of meteorological and air pollutant-specific monitoring equipment. DEQ operates most of the monitors while several tribes operate monitors on tribal lands. Data from the network are sent directly to engineers and scientists through a telemetry network.

Table 2 presents a summary of the monitoring stations used and parameters monitored during 2011. Some parameters were monitored for only part of the year.

Figure 1 shows a map of monitoring stations that were active in 2011. Monitoring stations are mainly located in high population areas; however, DEQ does monitor air quality in some rural areas. Some sites are selected to focus on the emissions of a single pollutant or group of sources (e.g., near a high-traffic volume or residential wood burning area). Monitor siting and monitoring objectives are discussed in the pollutant-specific sections of this report.

Criteria pollutants are measured using methods approved by EPA to assess Idaho's compliance with NAAQS. In addition, some pollutants of interest are measured using more than one method. These additional methods help engineers and scientists to better understand the presence and behavior of these pollutants.

Table 3 lists the methods used for the various pollutants. A variety of methods allow for continuous monitoring, and most can be accessed through telemetry for instantaneous concentration readings. Additional information on measurement methods is available at EPA's website: <http://www.epa.gov/ttn/amtic/>.

In addition to the criteria air pollutants described in this report, urban air toxic compounds were monitored at a Nampa site from 2003 to 2005 and at five other sites in 2007. If resources become available, DEQ may resume air toxics monitoring in the future. For details on air toxics and chemical toxicity, visit the EPA website at [www.epa.gov/ttn/atw/index.html](http://www.epa.gov/ttn/atw/index.html).

**Table 2. Monitoring network for 2011.**

Site	Location	PM <sub>10</sub> TEOM	PM <sub>2.5</sub> FRM	PM <sub>2.5</sub> TEOM	PM <sub>2.5</sub> BAM	O <sub>3</sub>	SO <sub>2</sub>	NO <sub>2</sub>	NO <sub>y</sub>	CO
Boise	Idaho Transportation Department— 3311 W. State Street					●		●		
Boise	Fire Station #5—16th and Front Street	●								
Boise	Eastman Building—166 N. 9th Street									●
Boise	White Pine Elementary—401 E. Linden Street					●				
Coeur d'Alene	Lancaster Road			●		●		●		
Franklin	East 4800 South		●							
Garden Valley	946 Banks-Lowman Road			●						
Grangeville	US Forest Service Compound			●						
Idaho City	3851 Highway 21			●						
Idaho Falls	Hickory and Sycamore			●						
Ketchum	111 W. 8th Street			●						
Lewiston	Sunset Park—1200 29th Street			●						
McCall	US Forest Service—500 North Mission Street			●						
Meridian	St. Luke's—520 E. Eagle Road.		●	●		●	●	●	●	●
Moscow	1025 Plant Sciences Road			●						
Nampa	Fire Station—923 1st Street	●	●	●						
Pinehurst	Pinehurst School—106 Church Street	●	●	●						
Pocatello	Garrett and Gould	●		●						
Pocatello	Wastewater Treatment Plant— Batiste and Chubbuck						●			
Salmon	618 N. Saint Charles Street		●		●					
Sandpoint	1601 Ontario	●		●						
Soda Springs	P4/Monsanto—5 Mile Road						●			
St. Maries	9th and Center		●	●						
Twin Falls	1913 Addison Avenue East			●						

Notes: PM<sub>10</sub> TEOM—particulate matter 10 micrometers, tapered element oscillating microbalance, continuous federal equivalent method; PM<sub>2.5</sub> FRM—particulate matter 2.5 micrometers, federal reference method; PM<sub>2.5</sub> TEOM—particulate matter 2.5 micrometers, tapered element oscillating microbalance, continuous special purpose monitor; PM<sub>2.5</sub> TEOM—particulate matter 2.5 micrometers, tapered element oscillating microbalance, continuous federal equivalent method; PM<sub>2.5</sub> BAM—particulate matter 2.5 micrometers, beta attenuation monitor, continuous federal equivalent method; O<sub>3</sub>—ozone, seasonal (May–September); SO<sub>2</sub>—sulfur dioxide; NO<sub>2</sub>—nitrogen dioxide, seasonal (May through September); NO<sub>y</sub>—total reactive nitrogen; CO—carbon monoxide; ●—trace




Figure 1. 2011 Idaho ambient air monitoring network.

**Table 3. Monitoring methods used in Idaho in 2011.**

Pollutant Code	Measurement	Method	Units
CO	Carbon monoxide	Gas filter correlation CO analyzer	Parts per million
NO <sub>x</sub> /NO <sub>y</sub>	Nitrogen oxides	Gas phase chemiluminescence	Parts per billion
O <sub>3</sub>	Ozone	Ultraviolet absorption	Parts per million
PM <sub>10</sub> TEOM (FEM)	PM <sub>10</sub> TEOM	TEOM gravimetric	Micrograms per cubic meter
PM <sub>2.5</sub> FRM	PM <sub>2.5</sub> reference	Gravimetric	Micrograms per cubic meter
PM <sub>2.5</sub> BAM 1020 (FEM)	PM <sub>2.5</sub> BAM	Beta attenuation	Micrograms per cubic meter
PM <sub>2.5</sub> TEOM FDMS (FEM)	PM <sub>2.5</sub> TEOM	TEOM gravimetric—Filter Dynamics Measurement System	Micrograms per cubic meter
PM <sub>2.5</sub> TEOM (SPM)	PM <sub>2.5</sub> TEOM	TEOM gravimetric	Micrograms per cubic meter
SO <sub>2</sub>	Sulfur dioxide	Ultraviolet fluorescence	Parts per billion
		Pulsed fluorescence	Parts per billion

Notes: Tapered element oscillating microbalance (TEOM); federal equivalent method (FEM); federal reference method (FRM); beta attenuation monitor (BAM); special purpose monitor (SPM)

### 3.1 Pollutant Monitoring

Coarse particulate (PM<sub>10</sub>) and fine particulate (PM<sub>2.5</sub>) are measured using a variety of methods in Idaho. EPA considers the federal reference method (FRM) or the federal equivalent method (FEM) to be most accurate for determining PM<sub>10</sub> and PM<sub>2.5</sub> concentrations. The FRM involves pulling in air (at a given flow rate) and trapping particles of a certain size (PM<sub>10</sub> or PM<sub>2.5</sub>) on a preweighed filter. The filter is weighed again, and the resulting mass is divided by volume of air sampled (determined from flow rate and amount of time) to provide concentration. Particles on the filter can be chemically analyzed later for more information about the sources of particulate matter. Unfortunately, the FRM does not provide continuous or timely information.

EPA has designated the tapered element oscillating microbalance (TEOM) continuous method an FEM for PM<sub>10</sub>. For PM<sub>2.5</sub> DEQ uses a specific variation of the TEOM, TEOM-Filter Dynamics Measurement System (FDMS), at the Pinehurst monitoring site. This variation is designated as an equivalent method for PM<sub>2.5</sub>, but the other TEOMs, despite the PM<sub>10</sub>, are not designated as such. The beta attenuation monitor (BAM) 1020 is designated as an equivalent method for PM<sub>2.5</sub>. DEQ uses this method at its Salmon monitoring site. DEQ uses the TEOM and TEOM-FDMS continuous methods to provide more time-resolved data (i.e., hourly averages) and to assess and forecast air quality in real-time or near real-time. DEQ also uses a variety of FEM gas analyzers for collecting data to determine compliance to the NAAQS. Some of these analyzers are also used to provide air quality forecasts near real-time. Data collected by methods not designated FRM or FEM cannot be used to determine compliance to NAAQS.

---

## 4 Monitoring Results

### 4.1 Ozone

Ozone, typically a summertime air pollution problem, forms when pollutants from internal combustion engines and industrial sources (e.g., paints, solvents, and gas vapors) react with sunlight. These pollutants are called ozone precursors and include VOCs and nitrogen oxides. Ozone can also be directly emitted by industrial sources. Ozone levels are usually highest in the afternoon because of the intense sunlight, warm temperatures, and the time required for ozone to form. These levels are highly affected by weather. DEQ monitored ozone from May through September 2011, as this is the time period specified by EPA requirements and the most likely time that high ozone levels will be observed.

Ozone is considered beneficial in the upper atmosphere because it helps to protect the earth from the sun's rays; however, ozone formed at ground level is unhealthy. Elevated concentrations of ground-level ozone can cause reduced lung function and respiratory irritation and can aggravate asthma. Ozone has also been linked to immune system effects ([www.epa.gov/ttn/oarpg/naaqsfin/o3health.html](http://www.epa.gov/ttn/oarpg/naaqsfin/o3health.html)). The damage ozone causes to the lungs typically heals within a few days, but repeated or prolonged exposure may cause permanent damage. People with respiratory conditions should limit outdoor exertion if ozone levels are high. Even healthy individuals may experience respiratory symptoms on a high ozone day. Ground-level ozone can also damage agricultural crops and forests, interfering with their ability to photosynthesize and grow.

Precursor chemicals that react with sunlight to produce ozone are generated primarily in large metropolitan areas. Because Idaho summers are normally hot and dry, ozone levels typically begin to rise in the late morning and peak in the late afternoon and early evening. This phenomenon follows closely with the time of day that the sun is the highest in the sky and temperatures are the hottest.

The ozone standard is defined so that the three highest ozone concentrations in any particular year can exceed the level of the standard while the area still maintains an *attainment* classification. However, if the 3-year averages of the 4th highest concentration exceed the level of the standard, the area is classified as *nonattainment* (Figure 2). Starting in 2008, the 3-year average (2006–2008) of the 4th highest 8-hour concentration will violate the NAAQS if it exceeds 0.075 ppm (0.076 ppm or higher).

Since 2002 DEQ has monitored ozone in Boise (Idaho Transportation Department, Whitney Elementary, and White Pine Elementary), Coeur d'Alene (Lancaster Road), Mayfield (Tilli Road), Middleton, and Meridian. Graphs presented in Figure 3–Figure 10 show trends in ozone levels at the monitoring stations in operation during 2011. For each station, the first graph presents daily maximum 8-hour average data for May through September. The shading on each graph corresponds to the AQI categories. The AQI categories of orange and above indicate NAAQS excursions. Breaks in the graphs are due to data being discarded as invalid. Data invalidation occurs when an instrument is taken off-line for routine maintenance, or there is a malfunction. Only valid data are shown on the graphs. The second graph presents the four highest concentrations observed during the year. The yellow circle presents the rolling 3-year

average. The 3-year average of the 4th highest concentration is the value used to assess compliance with the NAAQS.

Figure 11 shows a summary of the ozone monitoring data against the previous and new 8-hour federal standard. The figure shows that the state has remained at or below the previous ozone standard since monitoring began. The current ozone standard is closer to the 8-hour averages in the Treasure Valley, although the values have remained below the standard. For additional information on ozone, refer to Section 7, “Definitions” and Section 9, “Criteria Air Pollutants,” and visit <http://www.epa.gov/air/ozonepollution/>.





Figure 2. 2011 Idaho nonattainment and maintenance areas.

### Coeur d'Alene - Lancaster Rd. 8-Hour Ozone Daily maximum 8-hour concentration



\* Gaps in the charted data reflect times when valid data were not collected either from instrument malfunction, quality assurance failure, or equipment maintenance.

**Figure 3. Coeur d'Alene—Lancaster Road 8-hour ozone maximum concentration.**

### Ozone Measured at Coeur d'Alene - Lancaster Rd.

Four highest 8-hour concentrations  
and 3-year average of 4<sup>th</sup> highest





Figure 4. Coeur d'Alene—Lancaster Road highest 8-hour ozone concentrations and 3-year average of the 4th highest concentration.

### Meridian - St. Luke's 8-Hour Ozone Daily maximum 8-hour concentration



\* Gaps in the charted data reflect times when valid data were not collected either from instrument malfunction, quality assurance failure, or equipment maintenance.

**Figure 5. Meridian—St. Luke's 8-hour ozone daily maximum concentration.**

### Ozone Measured at Meridian - St. Luke's

Four highest 8-hour concentrations  
and 3-year average of 4<sup>th</sup> highest





Figure 6. Meridian—St. Luke's highest 8-hour ozone concentrations and 3-year average of the 4th highest concentration.



\* Gaps in the charted data reflect times when valid data were not collected either from instrument malfunction, quality assurance failure, or equipment maintenance.

**Figure 7. Boise—Idaho Transportation Department 8-hour ozone daily maximum concentration.**

### Ozone Measured at Boise - ITD

Four highest 8-hour concentrations  
and 3-year average of 4<sup>th</sup> highest




Figure 8. Boise—Idaho Transportation Department annual four highest 8-hour ozone concentrations and 3-year average of the 4th highest concentration.

### Boise - White Pine 8-Hour Ozone Daily maximum 8-hour concentration




Figure 9. Boise—White Pine 8-hour ozone daily maximum concentration.

### Ozone Measured at Boise - White Pine

Four highest 8-hour concentrations  
and 3-year average of 4<sup>th</sup> highest




Figure 10. Boise—White Pine highest 8-hour ozone concentrations and 3-year average of the 4th highest concentration.




Figure 11. Idaho 8-hour ozone concentrations and 3-year average of the 4th highest concentration.

## 4.2 Particulate Matter (10 micrometers)

Particulate matter includes solid matter and liquid droplets suspended in the air. Particles between 2.5 and 10 micrometers in diameter are called *coarse* particles.  $PM_{10}$  includes fine ( $PM_{2.5}$ ) and coarse particles. Coarse particles typically come from crushing or grinding operations and dust from roads.  $PM_{10}$  can aggravate respiratory conditions such as asthma. People with respiratory conditions should avoid outdoor exertion if  $PM_{10}$  levels are high.

The federal annual  $PM_{10}$  standard was revoked effective December 17, 2006, from a lack of evidence linking health problems to long-term exposure to coarse particle pollution. The 24-hour standard was not changed. EPA may choose to replace the  $PM_{10}$  standard in the future with a  $PM_{10-2.5}$  ( $PM_{coarse}$ ) standard, ranging from diameters 2.5 to 10 micrometers. Boise, Pocatello, Sandpoint, and Pinehurst have previously violated federal  $PM_{10}$  standards (Figure 2). Sandpoint and Pinehurst are currently nonattainment areas for  $PM_{10}$ . Pocatello (Portneuf Valley) and Boise (northern Ada County) were formerly nonattainment areas but are now considered to be maintenance areas for  $PM_{10}$ .

Idaho monitors  $PM_{10}$  using the continuous equivalent methods. The  $PM_{10}$  TEOM is a federal equivalent method. TEOM data are used to determine compliance to the  $PM_{10}$  NAAQS. The

TEOM method results are shown in Figure 12 and Figure 13. TEOM method data are also used to determine the daily AQI and to inform the public of air quality values in near real-time via DEQ’s web pages at <http://airquality.deq.idaho.gov/>.

The 3-year average daily maximum values shown in Figure 12 describe a variety of trends for the different monitor sites. The Boise and Pinehurst monitors registered decreased air quality values during the last year, while the Nampa showed increased values. Pocatello’s air quality values remained steady for the previous 3 years. Statistical summaries of the PM<sub>10</sub> concentrations are provided in Appendix A. The 24-hour PM<sub>10</sub> NAAQS is considered violated only if more than three exceedances occur during the consecutive 3-year period. For example, we could experience two exceedances in year one, none in year two, and one in year three and not violate the NAAQS.

Figure 13 demonstrates that Idaho’s airsheds, where monitoring is occurring, were in compliance with the daily NAAQS for PM<sub>10</sub> in 2011. The graph shows the 3-year average estimated exceedances of the 24-hour primary standard. None of the PM<sub>10</sub> monitors have violated the standard during the past 10 years.

For additional information on PM<sub>10</sub>, refer to Section 7 “Definitions” and visit [www.epa.gov/oar/particlepollution/](http://www.epa.gov/oar/particlepollution/).




Figure 12. Three-year average of daily maximum PM<sub>10</sub>.




Figure 13. Three-year average estimated exceedances of the daily PM<sub>10</sub> standard.

### 4.3 Particulate Matter (2.5 micrometers)

Particles 2.5 micrometers in diameter or less are called *fine* particles, or PM<sub>2.5</sub>. DEQ considers PM<sub>2.5</sub> to be one of the major air pollution concerns affecting a number of airsheds in Idaho. PM<sub>2.5</sub> generally comes from wood and agricultural burning, industrial boilers, and vehicle exhaust, including cars, diesel trucks, and buses. Fine particulate matter can also be formed secondarily in the atmosphere by chemical reactions of pollutant gases.

Exposure to PM<sub>2.5</sub> can have serious health effects. Fine particles are closely associated with increased respiratory disease, decreased lung function, and even premature death. Children, older adults, and people with some illnesses are more sensitive and more likely to develop heart or lung problems associated with PM<sub>2.5</sub>. People with respiratory or heart disease, older adults, and children should avoid outdoor exertion if PM<sub>2.5</sub> levels are high. PM<sub>2.5</sub> also significantly affects visibility.

DEQ measures PM<sub>2.5</sub> using three different methods—FRM, TEOM, and BAM. FRM was originally approved by EPA to determine PM<sub>2.5</sub> NAAQS compliance. This method involves pulling air through a size-selective inlet and a preweighed filter at a given flow rate, which traps particles of a certain size (in this case PM<sub>2.5</sub>) on the preweighed filter. The filter is weighed

again, and the net weight is divided by the volume of sampled air (determined from flow rate and amount of time) to provide the concentration. Unfortunately, FRM does not provide continuous or timely information. Idaho uses the TEOM method to provide more time-relevant data. The TEOM method accounts for the measurement of mass, via an oscillating filter element, to determine particulate matter present. The BAM method also provides data in near real-time. Its measurement method relies on the principle of attenuation to determine particulate concentrations. A final method of PM<sub>2.5</sub> measurement is achieved using a nephelometer monitor, which measures particulate matter using light scatter technology. This method is used during the agricultural burning season.

EPA provides FRM and FEM designation to monitoring methods that meet certain requirements. The designation allows the methods to be recognized by EPA as appropriate for NAAQS compliance determinations. The graphs in this section use data collected primarily from FRMs. The continuous data are primarily from TEOM methods, which typically are not designated as FRM or FEM but as special purpose monitors. An exception in DEQ's network—one TEOM method is designated an FEM—is the FDMS TEOM monitor located at Pinehurst. BAM, located at Salmon, is also an FEM. To improve the accuracy of the TEOM special purpose monitors, they are compared to the FRM values for a period of time, and calculations are made to determine the degree of difference between the two monitoring methods. The calculations are then applied to the current TEOM values in an attempt to make them *reference method-like*. Data gathered by the TEOM special purpose or nephelometer monitors cannot be used for NAAQS compliance determinations because they do not meet EPA equivalency requirements. States can request approval to use non-FRM and non-FEM monitors for NAAQS compliance through the approved regional method process. DEQ has not begun this process.

Figure 14 shows the 2011 3-year average of the 98th percentile 24-hour (daily) averages at Idaho monitoring stations against the federal standard. Franklin has violated the daily standard for the past 2 years, including 2011. Pinehurst has done the same. Salmon also violated the standard during 2011. These sites remain designated in attainment until a SIP determination is made by EPA for the 24-hour NAAQS, except for Franklin, which was included in the Cache Valley nonattainment area in 2010.

Figure 15 shows the 3-year average of the annual averages at each monitoring station against the federal standard. The data show that the annual standard of 15 µg/m<sup>3</sup> was not violated at any monitoring station.

Figure 16–Figure 33 show daily PM<sub>2.5</sub> concentrations measured at Idaho sites during 2011 using the continuous analyzers against a backdrop of AQI breakpoints. Most sites reflect the sustained nature of Good air quality throughout the state, with a few Yellow days interspersed. Five sites show significant spikes in the orange or Unhealthy for Sensitive Groups category. The excursions are likely the result of smoke from forest fires, residential woodburning, or windblown dust. A few of the graphs show some blank periods with no concentrations. These times indicate when a TEOM monitor was not functioning due to mechanical malfunctions or maintenance.

For additional information on particulate matter, refer to Section 7 “Definitions” and Section 9 “Criteria Air Pollutants,” and visit [www.epa.gov/oar/particlepollution/](http://www.epa.gov/oar/particlepollution/).




Figure 14. Three-year average 98th percentile daily PM<sub>2.5</sub> concentration (monitors operated in 2011).




Figure 15. PM<sub>2.5</sub> 3-year average annual mean (monitors operated in 2011).




Figure 16. Nampa—Fire Station PM<sub>2.5</sub> daily averages from continuous analyzer.

### Coeur d'Alene - Lancaster Rd. PM<sub>2.5</sub> daily averages from continuous analyzers




Figure 17. Coeur d'Alene—Lancaster Road PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 18. St. Maries PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 19. Sandpoint PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 20. Lewiston PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 21. Moscow PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 22. Twin Falls PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 23. Pocatello—Garrett and Gould PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 24. Idaho Falls PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 25. Garden Valley PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 26. Grangeville PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 27. Idaho City PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 28. Ketchum PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 29. McCall PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 30. Meridian—St. Luke's PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 31. Franklin PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 32. Pinehurst PM<sub>2.5</sub> daily averages from continuous analyzer.




Figure 33. Salmon PM<sub>2.5</sub> daily averages from continuous analyzer.

## 4.4 Carbon Monoxide

Carbon monoxide is an odorless, colorless gas that can enter the bloodstream through the lungs and reduce the amount of oxygen reaching organs and tissues. Carbon monoxide forms when the carbon in fuels does not burn completely. The majority of carbon monoxide comes from vehicle exhaust. In cities, 85%–95% of all carbon monoxide emissions come from motor vehicle exhaust.

Elevated levels of carbon monoxide in the ambient air can occur in urban canyon areas with heavy traffic congestion. The highest levels of carbon monoxide in the outside air typically occur during the colder months of the year when temperature inversions are more frequent. People with cardiovascular disease or respiratory problems might experience chest pain and increased cardiovascular symptoms, particularly while exercising, if carbon monoxide levels are high. High levels of carbon monoxide can affect alertness and vision even in healthy individuals.

Carbon monoxide monitoring stations are generally located in urban canyon areas with heavy traffic congestion, including central business areas, roadsides, and shopping malls. Idaho currently monitors carbon monoxide in Boise as a condition of EPA's *Northern Ada County (Boise), Idaho CO Maintenance Plan*. In 2009, trace carbon monoxide monitoring began at the NCore site in Meridian. Trace monitoring provides the ability to determine whether variations in observed concentrations below 1 ppm are from actual changes in atmospheric concentration or from poor sensitivity of older instruments at those low levels.

Figure 34 shows the second highest 8-hour concentrations at Idaho's monitoring sites versus the NAAQS from 2002 through 2011. The 2nd-highest concentration is displayed on these graphs because, under the federal rule, the 8-hour standard cannot be exceeded more than once per year (thus, choosing the 2nd highest). The data in these graphs demonstrate the sustained low concentrations that have been measured over the last 10 years. None of the 8-hour concentrations measured at any sites exceeded the NAAQS (9 ppm). The maximum 8-hour concentration for carbon monoxide in 2011 was 4.5 ppm, well below the 8-hour standard. These data are provided in Appendix A.

The NAAQS also includes a 1-hour standard for carbon monoxide of 35 ppm (cannot be exceeded more than once in any year). Measured 1-hour concentrations in Idaho are historically much lower than the 35 ppm standard; therefore, 1-hour carbon monoxide trends are not graphed. The maximum and 2nd-highest measured 1-hour carbon monoxide concentrations in 2011 were 20.4 ppm and 8.7 ppm, respectively. Additional 1-hour average carbon monoxide data are provided in Appendix A.

For additional information on carbon monoxide, refer to Section 7 "Definitions" and Section 9 "Criteria Air Pollutants," and visit <http://www.epa.gov/airquality/carbonmonoxide/>.




Figure 34. Carbon monoxide 2nd highest 8-hour concentration.

## 4.5 Sulfur Dioxide

Sulfur dioxide is a colorless, reactive gas produced by burning fuels containing sulfur, such as coal and oil, and by industrial processes. Historically, the greatest sources of sulfur dioxide were industrial facilities that derived their products from raw materials like metallic ore, coal, and crude oil, or that burned coal or oil to produce process heat (petroleum refineries, cement manufacturing, and metal processing facilities). Currently, on-road vehicles, marine craft, and diesel construction equipment also release significant sulfur dioxide emissions to the air.

People with asthma who are active outdoors may experience bronchoconstriction, where symptoms include wheezing, shortness of breath, and tightening of the chest. People should limit outdoor exertion if sulfur dioxide levels are high.

DEQ performs *hotspot* monitoring at the Pocatello and Soda Springs sites. Hotspot monitoring investigates pollution sources on a local scale. This monitoring assesses impacts from discreet sources to ambient air, rather than emissions monitored directly from a stack or chimney. In 2009, DEQ began trace sulfur dioxide monitoring at the NCore site in Meridian. Trace monitoring can determine whether variations in observed concentrations below 0.05 ppm are from actual changes in atmospheric concentration or from poor sensitivity of older instruments at

those low levels. On June 22, 2010, EPA revoked the annual and 24-hour standards and implemented a new 1-hour primary standard. The level was set at 75 ppb, and the level cannot exceed the 99th percentile of 1-hour daily maximum concentrations, averaged over 3 years. Three years of data for the averaging period will not be achieved until 2013, so no data are shown here. Table A-8 in Appendix A gives the first and second maximum and 99th percentile values recorded at the Meridian, Pocatello, and Soda Springs monitors in 2011.

Additional sulfur dioxide data are located in Appendix A. For information on sulfur dioxide, refer to Section 7 “Definitions” and Section 9 “Criteria Air Pollutants,” and visit <http://www.epa.gov/air/sulfurdioxide/>.

## 4.6 Lead

Lead is a highly toxic metal used for many years in household products, automobile fuel, and industrial chemicals. Airborne lead was associated primarily with automobile exhaust and lead smelters. The large reductions in lead emissions from motor vehicles have resulted in great reductions of ambient lead levels across the United States. Industrial processes, particularly primary and secondary lead smelters and battery manufacturers, are now responsible for most of the lead emissions.

People, animals, and fish are mainly exposed to lead by breathing and ingesting it in food, water, soil, or dust. Lead accumulates in the blood, bones, muscles, and fat. Infants and young children are especially sensitive to even low levels of lead. Lead can have health effects ranging from behavioral problems and learning disabilities to seizures and death.

According to EPA, the primary sources of lead exposure are lead-based paint, lead-contaminated dust, and lead-contaminated residual soils. Refer to EPA’s website for ways to limit your exposure to these lead sources: [www.epa.gov/ttnatw01/hlthef/lead.html](http://www.epa.gov/ttnatw01/hlthef/lead.html).

Lead has not been monitored in Idaho since 2002. With the phase-out of lead in fuel and the closure of the Bunker Hill lead smelter in Kellogg, airborne lead is no longer considered a public health concern in Idaho.

On November 12, 2008, EPA substantially strengthened the NAAQS for lead. EPA revised the level of the primary (health-based) standard from 1.5  $\mu\text{g}/\text{m}^3$  to 0.15  $\mu\text{g}/\text{m}^3$  and revised the secondary (welfare-based) standard to be identical in all respects to the primary standard. This level cannot be exceeded during any rolling 3-month average. In conjunction with strengthening the lead NAAQS, EPA promulgated new monitoring requirements. Monitoring is now required near lead sources that may contribute to violations of the lead NAAQS. Source-oriented monitoring is required near any source that emits more than 0.5 tons per year. Idaho does not have any sources of lead that trigger source-oriented monitoring. The monitoring regulations also require nonsource-oriented monitoring in metropolitan areas exceeding a 500,000 population at NCore multipollutant monitoring sites, beginning January 2012. DEQ will initiate lead monitoring at its NCore site in 2012.

For additional information on lead, refer to Section 7 “Definitions” and Section 9 “Criteria Air Pollutants,” and visit [www.epa.gov/air/lead/](http://www.epa.gov/air/lead/).

## 4.7 Nitrogen Dioxide

Nitrogen dioxide is a reddish brown, highly reactive gas that forms from the reaction of nitrogen oxide and oxygen in the atmosphere. The term NO<sub>x</sub>, which is frequently used, refers to both nitrogen oxide and nitrogen dioxide. Nitrogen dioxide will react with VOCs and can result in ozone. On-road vehicles like trucks and automobiles are the major sources of NO<sub>x</sub> in many airsheds. Industrial boilers and processes, home heaters, and gas stoves can also produce NO<sub>x</sub>. Nitrogen dioxide pollution is greatest during the cold weather seasons.

Nitrogen dioxide can cause respiratory symptoms such as coughing, wheezing, and shortness of breath in people with respiratory diseases such as asthma. Long-term exposure can lead to respiratory infections.

Motor vehicle manufacturers have been required to reduce NO<sub>x</sub> emissions from cars and trucks since the 1970s. NO<sub>x</sub> is not considered a significant pollution problem in Idaho. In 2011, DEQ operated one nitrogen dioxide monitor at its NCore site in Meridian. The monitoring objective was to assess ambient NO<sub>x</sub> concentrations for evaluating ozone formation processes during the ozone season.

The maximum 1-hour average of nitrogen dioxide measured in 2011 was 0.036 ppm. The averages observed have consistently been well below the hourly and annual NAAQS, as shown in Figure 35 and in the data in Appendix A. Until 2009, these averages could not be used to assess NAAQS compliance since the monitors were not operated for the entire year. Beginning in 2009, DEQ began monitoring nitrogen dioxide year-round at the NCore site in Meridian.

For additional information on nitrogen dioxide, refer to Section 7 "Definitions" and Section 9 "Criteria Air Pollutants," and visit <http://www.epa.gov/air/nitrogenoxides/>.

### Idaho Nitrogen Dioxide (NO<sub>2</sub>) Annual 1-hour average vs. standard




Figure 35. Nitrogen dioxide annual 1-hour average.

## 5 Air Quality Index

The AQI is reported according to a 500-point scale for each of the major criteria air pollutants: ozone, particulate matter (PM<sub>10</sub> and PM<sub>2.5</sub>), carbon monoxide, nitrogen dioxide, and sulfur dioxide. The *worst denominator* determines the ranking. For example, if an area has a carbon monoxide value of 132 on a given day and all other pollutants are below 50, the AQI for that day would be 132. The AQI scale breaks down into six categories. Each category has a corresponding color, shown in Table 4. For information on the concentration breakpoints for each pollutant, refer to Table A-1 in Appendix A.

**Table 4. US Environmental Protection Agency Air Quality Index breakpoint definitions.**

Levels of Health Concern	Numeric Value	Meaning
Good	0–50	Air quality is satisfactory, and air pollution poses little or no risk.
Moderate	51–100	Air quality is acceptable; however, for some pollutants there may be a moderate health concern for a small number of people who are unusually sensitive to air pollution.
Unhealthy for sensitive groups	101–150	Members of sensitive groups may experience health effects. The general public is not likely to be affected.
Unhealthy	151–200	Everyone may begin to experience health effects. Members of sensitive groups may experience more serious health effects.
Very unhealthy	201–300	Health alert: everyone may experience more serious health effects.
Hazardous	301–500	Health warnings of emergency conditions. The entire population is more likely to be affected.

The AQI is a national index, so the values and colors used to show local air quality and the associated level of health concern will be the same everywhere in the United States. The number of Good air quality days continues to dominate regionally in Idaho. However, periods occurred when the air quality degraded into Moderate, Unhealthy for Sensitive Groups, and Unhealthy categories. Table 5 shows the number of days in each AQI category in Idaho counties where air quality is monitored. In 2011, the highest AQI value of 166 was recorded in Franklin County. This value was in the Unhealthy range.

In Figure 36–Figure 52, the AQI graphs present the distribution of air quality for each individual county. The AQI data summaries for each county, which support the graphs' data, are located in Table A-2 in Appendix A.

Table 5. 2011 Air Quality Index yearly summary.

2011 AQI Ratings						
2011		Number of days in AQI category				
County	Total number of AQI days	Good	Moderate	Unhealthy for sensitive groups	Unhealthy	Highest AQI
Ada	365	315	47	3	—	115
Bannock	365	247	115	3	—	114
Benewah	361	320	41	—	—	99
Bonner	340	340	—	—	—	37
Butte	364	348	16	—	—	77
Canyon	358	336	21	1	—	110
Caribou	365	353	12	—	—	96
Cassia	195	192	3	—	—	58
Custer	91	89	2	—	—	75
Franklin	312	254	49	4	5	166
Idaho	306	306	—	—	—	50
Kootenai	183	182	1	—	—	51
Latah	365	354	11	—	—	70
Lemhi	359	222	125	11	1	152
Nez Perce	365	345	20	—	—	68
Shoshone	360	244	98	18	—	147
Twin Falls	350	335	15	—	—	66




Figure 36. Air quality for Ada County.




Figure 37. Air quality for Bannock County.




Figure 38. Air quality for Benewah County.




Figure 39. Air quality for Bonner County.




Figure 40. Air quality for Butte County.




Figure 41. Air quality for Canyon County.




Figure 42. Air quality for Caribou County.




Figure 43. Air quality for Cassia County.




Figure 44. Air quality for Custer County.




Figure 45. Air quality for Franklin County.




Figure 46. Air quality for Idaho County.




Figure 47. Air quality for Kootenai County.




Figure 48. Air quality for Latah County.




Figure 49. Air quality for Lemhi County.




Figure 50. Air quality for Nez Perce County.




Figure 51. Air quality for Shoshone County.




Figure 52. Air quality for Twin Falls County.

## 6 Impaired Air Quality

### 6.1 Winter Burn Bans

Idaho has a winter-impaired air quality program primarily targeting sources of particulate matter from open outdoor burning, prescribed fire use, and woodstoves and fireplaces. Idaho's program is implemented through local ordinances in those areas with winter inversion problems. These ordinances specify that public actions such as open burning bans or indoor wood burning bans take place whenever DEQ reports an AQI above a certain value and air stagnation conditions are forecasted to continue for at least 24 hours.

The DEQ web page, [Daily Air Quality Reports and Forecasts](#), lists the daily air quality in many cities and regions in Idaho. Each report lists the air pollutant being monitored, AQI, and burn restrictions, if any, for the day. Residents interested in air quality information can visit the web page to see the forecast for their area.

DEQ issues an air quality advisory for specific locations between 2:00 p.m. and 4:00 p.m. for the next day when air quality is forecast to be poor. The advisories issued are based upon the expected conditions for the next day. Updates are sent out on weekend days at the same times if conditions are expected to be poor. These advisories are provided to local media outlets and to others through e-mail notification lists.

### 6.2 Summer Ozone Alerts

DEQ forecasts pollution conditions for ozone in the Treasure Valley and Kootenai County using pollutant monitoring data and meteorological information. Because ozone needs heat and sunlight to form, it is considered a summertime problem and is only monitored from May 1 through September 30. Ozone pollution can rise to high levels when the valley experiences hot days with few clouds in the sky. The Treasure Valley and Kootenai County tend to see daily ozone levels that begin to rise in the late morning and peak in the late afternoon and early evening. This phenomenon follows closely with the time of day that the sun is the highest in the sky through the time temperatures are the hottest. Since we have no control over our weather characteristics, we have to focus on controlling what we put into our air. Under Yellow or Moderate alerts, the public is requested to change certain behaviors to prevent further deterioration of air quality. These alerts are reported to local media outlets and to others through an e-mail notification list.

## 7 Definitions

### Air Toxics

Air toxics are broadly defined as almost 700 pollutants that DEQ considers to be potentially harmful to human health and the environment. These pollutants are listed in the “Rules for the Control of Air Pollution in Idaho” (IDAPA 58.01.01.585 and 586) at <http://adminrules.idaho.gov/rules/current/58/0101.pdf>. Hazardous air pollutants (see below) are included in this list to identify them as a subset of air toxics.

### Criteria Air Pollutant

The Clean Air Act of 1970 defined six criteria air pollutants and established ambient concentrations of each to protect public health. EPA periodically revises the original concentration limits and methods of measurement, most recently in 2010. Table 1 provides the list and the allowed ambient concentrations.

### Hazardous Air Pollutant

A hazardous air pollutant (HAP) is an air contaminant identified as toxic in the federal Clean Air Act, §112(b). Currently listed, 188 pollutants are considered HAPs: <http://www.epa.gov/ttn/atw/188polls.html>.

### Temperature Inversions

The earth gains and loses most of its energy at its surface. It is warmed by solar heating during the day and cooled by radiation emissions at night. During the late morning and afternoon hours, the air near the surface is warmer than the air aloft and allows for good pollutant dispersion (vertical mixing may be 1,500 meters or more). At night with clear skies, the surface radiates heat into outer space, creating cooler air at the surface and warmer air aloft. Warmer air above cooler air (temperature inversion) is a stable condition and limits the upward movement of pollution because the warmer air acts as a barrier. With little or no wind, pollutants are trapped near the surface (vertical mixing may be 200 meters or less) and can reach high levels of concentration.

### Volatile Organic Compound

A volatile organic compound (VOC) is a gas emitted from certain solids and liquids that participates in atmospheric photochemical reactions. This excludes all compounds determined to have negligible photochemical reactivity by EPA and listed in 40 CFR 51.100(s) in effect July 1, 1998.

### Visibility/Regional Haze

Visibility is often explained in terms of visual range and light extinction. Visual range is the maximum distance—usually miles or kilometers—that you can see a black object against the horizon. Light extinction is the sum of light scattering and light absorption by fine particles and gases in the atmosphere. The more light extinction, the shorter your visual range will be.

Reduced visibility (or visual range) is caused by weather (clouds, fog, and rain) and air pollution (fine particles and gases). The major pollution contributor to reduced visibility is fine particulate matter (PM<sub>2.5</sub>) emissions, which are transported aloft and may remain suspended for a week or longer. Fine particles have a greater impact than coarse particles at locations far from the emitting source because they remain suspended in the atmosphere longer and travel farther. PM<sub>2.5</sub> also presents some of the most serious health hazards to the public, so you can roughly assume that the worse the visibility, the unhealthier the air is to breathe.

## **8 Pollution Sources**

### **Area Sources**

Area sources are categories of pollution sources, in which each individual industrial source emits pollutants below the thresholds for a point-source facility designation, and includes other categories that are a result of human activities. Area sources are best estimated at a county level in association with population numbers (e.g., natural gas use for home heating, gas stoves, or woodstoves).

### **Biogenics**

Biogenics are natural sources such as trees, plants, grass, crops, and soils. The worldwide emissions rate of these natural hydrocarbons has been estimated to exceed that of nonmethane hydrocarbons originating from human sources. Isoprene, one of the major constituents of biogenic emissions, is very photoreactive and makes biogenic VOCs, a contributor in ozone formation.

### **Emission Factor**

Emission factor is a value derived from source tests, material balance calculations, or engineering comparisons with similar processes. It is used to estimate emissions from process quantities.

### **Nonroad Mobile Sources**

Nonroad mobile sources include farm vehicles, on-site construction/industrial vehicles, logging equipment, small marine craft, aircraft, trains, lawn and garden equipment, and off-road trail machines.

### **On-road Mobile Sources**

On-road mobile sources include cars, trucks, sport utility vehicles, motorcycles, and buses.

### **Point Sources**

For the every-third-year statewide emissions inventory, point sources are defined as facilities that have actual annual air pollutant emissions equal to, or exceeding, 1,000 tpy of carbon monoxide; 100 tpy of NO<sub>x</sub>, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>x</sub>, or VOCs; or 5 tpy of lead.

## Registered Facility

The total of all pollutant-emitting activities located on adjacent or contiguous properties owned or operated by one person or a corporate entity. It includes all of the pollutant-emitting buildings, processes, structures, equipment, control apparatuses, and storage areas at a facility.

## 9 Criteria Air Pollutants

### Ozone

- **What is it?**  
Ozone (O<sub>3</sub>), a bluish-colored gas molecule with a strong odor, is composed of three atoms of oxygen. In the upper atmosphere, ozone occurs naturally and partially absorbs the sun's harmful ultraviolet rays. Ozone at ground level is a summertime air pollution problem.
- **How is it caused?**  
Ozone forms when photochemical pollutants from cars, trucks, and industrial sources react with sunlight. Ozone-forming pollutants include NO<sub>x</sub> and VOCs; even gasoline-powered yard equipment, paints, solvents, and off-road vehicle motors contribute.
- **When does it happen?**  
Ozone pollution is most common in the summer months, when sunlight and stable atmospheric conditions occur. Ozone levels are usually highest in the afternoon, as sunlight photochemically transforms NO<sub>x</sub> and VOCs into ozone.
- **Who is affected?**  
Adults and children, who are active outdoors, people with respiratory disease such as asthma, and people with unusual sensitivity to ozone. During physical activity, ozone penetrates deeper into the lungs and can do more damage.

Ozone is a reactive gas. For this reason, high ozone concentrations can cause respiratory distress and disease in humans, decreased yields of agricultural crops and forests, and damage to some rubber products, plastics, and paints used outdoors. National crop losses from ozone exposure are estimated at \$3 billion to \$5 billion annually. Forest losses are harder to estimate.

- **What are the health effects?**  
Ozone can cause coughing and throat irritation, make deep vigorous breathing more difficult, and increase the chance of respiratory infections. It increases sensitivity to allergens and can trigger asthma attacks. The damage it causes to the lungs heals within a few days, but repeated or prolonged exposure may cause permanent damage.
- **What can I do about it?**  
If ozone levels are high and you have a respiratory condition or are normally active outdoors, try to limit your outdoor exertion.

In the United States, management of ozone and other photochemical oxidants has been a

major goal of federal and state clean air legislation (Clean Air Act of 1970). Although many of the pollution control efforts required by the CAA have been implemented, efforts to decrease ozone pollution have been only partially successful.

- **Where is it measured?**

Unlike other pollutants monitored here in Idaho, ozone is formed when precursor compounds react in the atmosphere. Winds transport ozone and precursor emissions from one area to another. For the Treasure Valley, ozone precursors are emitted into the air in urban areas of the airshed and subsequently travel southeasterly to more rural areas as they react to form ozone. As a result, for the Treasure Valley airshed, DEQ has monitors in various locations. Another ozone monitor has been running in the Coeur d'Alene area since 2005.

## Particulate Matter

- **What is it?**

Particulate matter (PM) includes both solid matter and liquid droplets suspended in the air. Particles smaller than 2.5 micrometers in diameter are called *fine* particles, or  $PM_{2.5}$ . Particles between 2.5 and 10 micrometers in diameter are called *coarse* particles.  $PM_{10}$  includes both fine and coarse particles. DEQ considers  $PM_{2.5}$  to be one of the major air pollution concerns affecting our state.

- **How is it caused?**

$PM_{2.5}$  comes from all types of combustion, including cars, diesel trucks, power plants, wood burning, and from some industrial processes. It can also be formed in the atmosphere by chemical reactions of pollutant gases. The *coarse* particles in  $PM_{10}$  typically come from crushing or grinding operations and dust from roads.

- **When does it happen?**

Daily  $PM_{2.5}$  trends in urbanized areas suggest that  $PM_{2.5}$  levels peak in association with traffic flow and rush hour periods. Periods of stagnate weather patterns, such as when surface inversions typically occur, contribute to elevated  $PM_{2.5}$  trends.

- **Who is affected?**

People with asthma and heart or lung disease, the elderly, and children are affected.  $PM_{2.5}$  also significantly affects visibility.

- **What are the health effects?**

Fine particulates ( $PM_{2.5}$ ) pose a greater risk to human health than coarse particulates, because they penetrate deeper into the respiratory system.  $PM_{2.5}$  exposure can have serious health effects. People with heart or lung diseases are at increased risk of attacks or premature death. Children and the elderly are more likely to develop heart or lung problems.  $PM_{10}$  can aggravate respiratory conditions such as asthma.

**What can I do about it?**

If PM<sub>2.5</sub> levels are high, people with respiratory or heart disease, the elderly, and children should avoid outdoor exertion. If PM<sub>10</sub> levels are high, people with respiratory conditions should avoid outdoor exertion.

- **Where is it measured?**

Due to the health risks associated with particulate matter, both PM<sub>2.5</sub> and PM<sub>10</sub> are monitored in various population-oriented locations throughout Idaho.

## **Carbon Monoxide**

- **What is it?**

Carbon monoxide (CO) is an odorless, colorless gas that can enter the bloodstream through the lungs and reduce the amount of oxygen that reaches organs and tissues.

- **How is it caused?**

Carbon monoxide forms when the carbon in fuels does not burn completely. Vehicle exhaust contributes 60% of all carbon monoxide. In cities, that contribution can be as high as 95%.

- **When does it happen?**

Carbon monoxide pollution is at its worst in cold weather because fuels burn less efficiently in low temperatures. Carbon monoxide levels usually peak during morning and evening rush hours.

- **Who is affected?**

People with cardiovascular disease, such as angina, or cardiovascular or respiratory problems, also fetuses and young infants.

- **What are the health effects?**

Chest pain and increased cardiovascular symptoms, particularly while exercising. High levels of carbon monoxide can even affect alertness and vision in healthy individuals.

- **What can I do about it?**

If carbon monoxide levels are high, limit exertion and avoid sources of carbon monoxide such as heavy traffic.

- **Where is it measured?**

Carbon monoxide monitoring stations are located in urban canyon areas with heavy traffic congestion. These include central business areas, roadsides, and shopping malls. The Boise carbon monoxide monitor is located in downtown Boise and monitors carbon monoxide as part of an air quality maintenance plan. Beginning in 2009, carbon monoxide is also monitored in Meridian.

## Sulfur Dioxide

- **What is it?**  
Sulfur dioxide (SO<sub>2</sub>) is a colorless, reactive gas.
- **How is it caused?**  
Sulfur dioxide is produced by burning sulfur-containing fuels such as coal and oil and by some industrial processes.
- **Where does it happen?**  
The highest concentrations of sulfur dioxide are usually near large industrial sources.
- **Who is affected?**  
People with asthma who are active outdoors.
- **What are the health effects?**  
Bronchoconstriction can cause wheezing, shortness of breath, and tightening of the chest. When exposure to sulfur dioxide ends, the symptoms should clear up within an hour.
- **What can I do about it?**  
If sulfur dioxide levels are high, limit your outdoor exertion.
- **Where is it measured?**  
Because the large primary sources of sulfur dioxide in Idaho are industrial, DEQ monitors for sulfur dioxide near large facilities with high sulfur dioxide emissions. The monitors running in 2011 were in Pocatello, Soda Springs, and Meridian.

## Lead

- **What is it?**  
Lead (Pb) is a highly toxic metal that was used for many years in household products, automobile fuel, and industrial chemicals.
- **How is it caused?**  
Locally, airborne lead is associated primarily with automobile exhaust and lead smelters. Since the phase-out of lead in fuels, cars and trucks are no longer a significant source of lead. The Kellogg Bunker Hill Mine ceased operations in 1981, which also contributed to lead source reduction.
- **When does it happen?**  
Lead concentrations are likely to be highest near sources where current or former lead smelting/processing operations caused particle fallout, especially in nearby soils such as unpaved parking lots.
- **Who is affected?**  
Everyone. Children who are 6 years and younger are most at risk.
- **What are the health effects?**  
Lead can have health effects ranging from behavioral problems and learning disabilities to seizures and death.

- **What can I do about it?**  
According to EPA, the primary sources of lead exposure are lead-based paint, lead-contaminated dust, and lead-contaminated residual soils. Refer to EPA's website at <http://www.epa.gov/ttn/atw/hlthef/lead.html> for ways to limit your exposure to these lead sources.
- **Where is it measured?**  
Due to the phase-out of leaded fuels and the closure of Idaho's only lead smelter in 1981, DEQ discontinued monitoring for airborne lead. Historical monitoring was continued until 2002 but was discontinued due to the low levels being measured. With the lowering of the lead standard, DEQ will resume lead monitoring in 2012 at the NCore site in Meridian.

## Nitrogen Dioxide

- **What is it?**  
Nitrogen dioxide (NO<sub>2</sub>) is a reddish brown, highly reactive gas that forms from the reaction of nitrogen oxide (NO) and oxygen in the atmosphere. Nitrogen dioxide will react with VOCs and can result in ozone formation.
- **How is it caused?**  
High temperature combustion sources such as power plants and automobiles are major producers of nitrogen oxide. Home heaters and gas stoves can also produce nitrogen oxide.
- **When does it happen?**  
Nitrogen dioxide pollution is greatest in cold weather. It follows a similar trend to carbon monoxide.
- **Who is affected?**  
Children and people with respiratory diseases, such as asthma.
- **What are the health effects?**  
Nitrogen dioxide can cause respiratory symptoms such as coughing, wheezing, and shortness of breath. Long-term exposure can lead to respiratory infections.
- **What can I do about it?**  
Since the 1970s, motor vehicle manufacturers have been required to reduce nitrogen oxide emissions from cars and trucks. It is not a significant pollution problem in Idaho.
- **Where is it measured?**  
Nitrogen dioxide is not a major concern in Idaho. It was measured during 2011 at the NCore site in Meridian.

## Air Quality Data Summaries

Table A-1. Calculation and breakpoint for the Air Quality Index.

Breakpoints for Criteria Pollutants							AQI Categories	
O <sub>3</sub> (ppm) 8-hour	O <sub>3</sub> (ppm) 1-hour <sup>a</sup>	PM <sub>2.5</sub> (µg/m <sup>3</sup> ) 24-hour	PM <sub>10</sub> (µg/m <sup>3</sup> ) 24-hour	CO (ppm) 8-hour	SO <sub>2</sub> (ppb) 1-hour	NO <sub>2</sub> (ppb) 1-hour	AQI value	Category
0.000–0.059	—	0.0–15.4	0–54	0.0–4.4	0-35	0-53	0–50	Good
0.060–0.075	—	15.5–35.4	55–154	4.5–9.4	36-75	54-100	51–100	Moderate
0.076–0.095	0.125–0.164	35.5–65.4	155–254	9.5–12.4	76-185	101-360	101–150	Unhealthy for sensitive groups
0.096–0.115	0.165–0.204	65.5–150.4	255–354	12.5–15.4	186-304(c)	361-649	151–200	Unhealthy
0.116–0.374	0.205–0.404	150.5–250.4	355–424	15.5–30.4	305-604(c)	650-1249	201–300	Very unhealthy
(b)	0.405–0.504	250.5–350.4	425–504	30.5–40.4	605-804(c)	1250-1649	301–400	Hazardous
(b)	0.505–0.604	350.4–500.4	505–604	40.5–50.4	805-1004(c)	1650-2049	401–500	

- a. Areas are generally required to report the AQI based on 8-hour ozone values. However, there are a small number of areas where an AQI based on 1-hour ozone values would be safer. In these cases, in addition to calculating the 8-hour ozone value, the 1-hour ozone value may be calculated, and the greater of the two values reported.
- b. Eight-hour ozone values do not define higher AQI values ( $\geq 301$ ). AQI values of 301 or greater are calculated with 1-hour ozone concentrations.
- c. One-hour SO<sub>2</sub> values do not define higher AQI values ( $\geq 200$ ). AQI values of 200 or greater are calculated with 24-hour SO<sub>2</sub> concentrations.

Notes: ppm = parts per million; µg/m<sup>3</sup> = micrograms per cubic meter; ppm = parts per million; ppb = parts per billion

For more detailed information about the AQI and the pollutants it measures, go to <http://www.airnow.gov/>.

Table A-2. 2011 Air Quality Index summary report.

2011		Number of Days for AQI Categories					Max AQI	Number of Days for Main AQI Pollutants					
County	Total Number of AQI Days	Good	Moderate	Unhealthy for Sensitive Groups	Unhealthy	Very Unhealthy		CO	NO <sub>2</sub>	O <sub>3</sub>	SO <sub>2</sub>	PM <sub>2.5</sub>	PM <sub>10</sub>
Ada	365	315	47	3			115	11	9	231		49	65
Bannock	365	247	115	3			114				170	81	114
Benewah	361	320	41				99					361	
Bonner	340	340					37						340
Butte	364	348	16				77			361		3	
Canyon	358	336	21	1			110					55	303
Caribou	365	353	12				96				365		
Cassia	195	192	3				58			195			
Custer	91	89	2				75					91	
Franklin	312	254	49	4	5		166					312	
Idaho	306	306					50					306	
Kootenai	183	182	1				51			183			
Latah	365	354	11				70					365	
Lemhi	359	222	125	11	1		152					359	
Nez Perce	365	345	20				68					365	
Shoshone	360	244	98	18			147					345	15
Twin Falls	350	335	15				66					350	

Table A-3. 2011 monitor values summary for PM<sub>2.5</sub>.<sup>a</sup>

Data Year	County	# Obs	24-hour PM <sub>2.5</sub>							Annual Mean	Monitor #	Site ID	Site Address	City
			1st Max	2nd Max	3rd Max	4th Max	98th %	# Exceed						
2011	Ada	117	34.8	32.5	28.5	24.3	28.5	0	6.98	1	160010010	St. Luke's-520 S. Eagle Rd.	Meridian	
2011	Benewah	59	34.6	28.8	22.5	21.7	28.8	0	8.01	1	160090010	9th and Center	St. Maries	
2011	Canyon	61	29.7	22.5	21.6	21.0	22.5	0	8.92	1	160270002	Nampa Fire Station-923 1st St.	Nampa	
2011	Franklin	113	55.1	45.2	39.5	23.5	39.5	3	7.43	1	160410001	Water Treatment Facility-East 4800 South	Franklin	
2011	Lemhi	357	58.0	54.3	51.5	44.2	37.1	>10	12.21	3	160590004	618 N. St. Charles St.	Salmon	
2011	Shoshone	337	54.1	51.6	50.9	45.9	43.8	>10	11.96	4	160790017	Pinehurst School-106 Church St.	Pinehurst	

- a. Values indicated come from Federal Reference or Equivalent Method measurements.
- b. The # exceed indicates the number of times measurements exceeded the National Ambient Air Quality Standard (NAAQS) of 35.5 micrograms per cubic meter ( $\mu\text{g}/\text{m}^3$ ). Exceedances themselves do not cause an area to be designated nonattainment, but they can cause the 98th percentile value to be higher, which when averaged with the previous 2 years, can cause an area to be designated nonattainment. For example, in 2008, the 98th percentile for Pinehurst was above the standard at  $36.2 \mu\text{g}/\text{m}^3$ . Since the 3-year average of the 2006-2008 98th percentile is  $34.7 \mu\text{g}/\text{m}^3$ , which is below the standard, the area is classified as attainment.

Table A-4. 2011 monitor values summary for PM<sub>10</sub>.<sup>a</sup>

Data Year	County	24-hour PM <sub>10</sub>							Monitor #	Site ID	Site Address	City
		# Obs	1st Max	2nd Max	3rd Max	4th Max	# Exceed Actual	Estimated Exceedances				
2011	Ada	8,493	59	55	47	45	0	0	3	160010009	Fire Station #5– 16th and Front	Boise
2011	Bannock	8,567	72	69	65	62	0	0	3	160050015	Garret and Gould	Pocatello
2011	Bonner	8,077	40	36	32	31	0	0	3	160170005	1601 Ontario St.	Sandpoint
2011	Canyon	8,530	174	108	68	68	1	1	2	160270002	Nampa Fire Station–923 1st St.	Nampa
2011	Shoshone	8,256	39	37	37	36	0	0	3	160790017	Pinehurst School– 106 Church St.	Pinehurst

a. Values indicated come from Federal Reference or Equivalent Method measurements.

Table A-5. 2011 monitor values for ozone.

Data Year	County	8-hour O <sub>3</sub>							% Days	Monitor #	Site ID	Site Address	City
		1st Max	2nd Max	3rd Max	4th Max	Days > Std	Required Days	# Days					
2011	Ada	0.071	0.071	0.070	0.069	0	153	108	71	1	160010010	St. Luke's–520 S. Eagle Rd.	Meridian
2011	Ada	0.070	0.065	0.064	0.062	0	153	128	84	1	160010017	White Pine Elementary–401 E. Linden	Boise
2011	Ada	0.068	0.063	0.062	0.060	0	153	138	90	1	160010019	Idaho Transportation Dept.–3311 W. State St.	Boise
2011	Kootenai	0.060	0.059	0.058	0.058	0	153	150	98	1	160550003	Lancaster Rd.	Coeur d'Alene

Table A-6. 2011 monitor values summary for carbon monoxide.

Data Year	County	1-hour CO			8-hour CO			Monitor #	Site ID	Site Address	City	
		# Obs	1st Max	2nd Max	# Exceed	1st Max	2nd Max					# Exceed
2011	Ada	8,254	1.420	1.410	0	1.0	0.8	0	1	160010010	St. Luke's-520 S. Eagle Rd.	Meridian
2011	Ada	8,395	20.4	8.7	0	4.5	1.6	0	1	160010014	Eastman Building-166 N. 9th St.	Boise

Table A-7. 2011 monitor values summary for nitrogen dioxide.

Data Year	County	1-hour NO <sub>2</sub>			Annual NO <sub>2</sub>		Monitor #	Site ID	Site Address	City
		# Obs	1st Max	2nd Max	Mean	# Exceed				
2011	Ada	661	0.036	0.034	0.0105	0	1	160010010	St. Luke's-520 S. Eagle Rd.	Meridian

Table A-8. 2011 monitor values summary for sulfur dioxide.

Data Year	County	1-hour SO <sub>2</sub>			99th percentile	Site ID	Site Address	City
		# Obs	1st Max	2nd Max				
2011	Ada	8,225	0.0123	0.0111	0.0076	160010010	St. Luke's-520 S. Eagle Rd.	Meridian
2011	Bannock	7,025	0.1061	0.1024	0.0752	160050004	Sewage Treatment Plant-Batiste and Chubbuck	Pocatello
2011	Caribou	8,711	0.0727	0.0626	0.0532	160290031	P4/Monsanto-Five Mile Rd.	Soda Springs